Short Communication



ISSN: 2059-0377

THE Asterias rubens complement system: Comparisons with lower vertebrates

Michel Leclerc^{1*} and Nicolas Kresdorn²

¹556 rue Isabelle Romée, 45640 Sandillon, France ²GenXpro, Frankfurt, Germany

Abstract

Seven complement components have been discovered in 2013. Another component: the C6 component was found in sea star, when compared to rainbow trout genome.

Introduction

We have recently described the "Sea star complement Evidence" [1]. We remarked that C6 and C7 components were missing in sea star transcriptome when compared to mouse one.

An extensive study allowed us to research these components in less evolved animals (phylogenetically speaking) than mouse. Genomic features of the rainbow trout: *Oncorhynchus mykiss* have helped us, in this study.

At this point, we were attempting to determine how many similar complement components might be present in *Asterias rubens* (Invertebrate) and in *Oncorhynchus mykiss* (Vertebrate).

Materials and methods

Sea stars Asterias rubens were obtained from the Biology Institute (Gothenbugh University) Immunizations, genomic studies were already described [1]. After ligation of adapters for Illumina's GSII sequencing system, the cDNA was sequenced on the Illumina GSII platform sequencing.

1.100 bp from one side of the approximately 200 bp fragments. Sequences were assembled using Velvet [2].

Results

Three complement components: C1r, C4, C1 inhibitor of the classical activation pathway have been fully sequenced in rainbow trout [3].

C6 was discovered in trout in 2006 [4].

Sea star C1q subunits A, B, C, were sequenced in A. Rubens [1].

C2, C4B, and C3 which is central in mammals to both the classical and alternative pathways, C9,

C5, C8 were also sequenced (1) in Asterias rubens

As for C6, it was shown as following, when compared to Oncorhynchus mykiss genome:

One contig (Contig11285 m.9708) could be annotated via BLASTX

to Oncorhynchus mykiss "Complement component C6" from the Trembl database, with an e-value of 3.75e-13. On an aligned region of 113 amino acids, 37 positive and 56 identical amino acids were found.

Т С **G** А С А С Т Т А С А А А А А **G** С А Т С Т С А А С С С -G A G T A G G A A G G A A T C T C T T T T A G T T T A T T T T G T G C T C C C T T T G G T A T C A G T T T A G A T C C A C A C A A C C T G T G A A A A A C T T C A G T A C T -T A C T A G A T T T C G C C A A C G C A A C G G T A A A C G A G T C A T T T G A T T T T G A C C A T C A T C A A C T -GAAGCAACGCACGTAATACACAAAACGG A A C A T T T T G T G T G T G T A G T T T C C A G C G A T T C -G A G A A G C A A A T C A A A G A C A A G A T G T C T T T A C C C A G T G A T G T T G A A A C A G A C T C C G T C A T G -G A T A G T C C A G C A G A G A T T C A T A T G A A C A T G A A T A A G C T A C A A T C T A A A C T T C C C A G C G T -TACTCAAGACGAGAGATTTGACTCCGGAATTG A C T C G T T A C G T T C T G T T G A T T C G G C G T A C T -G C T T G A G C T T C G A A A G G G A A T C G A G C C T G G CTTCGATAAATGAGAAGACGTCTCTCACAT-C A C A C C T G C A A C A G C T C C A T C T T T C A C A T G G A C A G T A G A A G A C A T C G A T G A A G C T T A T C A T G A T G A G T G T A C T A T G T C T G A A A C A C T C -G A C A A T T T G G A A G A A A C T G C A A G A A T T G T G G A A T A T C C T G A A C A A A G A T G C A C G G G A C G T C T -T A C A G A T G A T G C C T T C G A C C A A G A C C A A G A G G G A G A T A C G C C C C T T C A T C T T G C T A T -T A T T C A T A A G G A A G T G G A C T T C G C A G A A A A A T T C A T C A T C T T T G T T G C A G A T C C T G A G T T A C T -G A A C A T C A G C A A T G A T C T T A T G C A G A C T C

Correspondence to: Michel Leclerc, 556 rue Isabelle Romée, 45640 Sandillon, France, **E-mail:** mleclerc45@gmail.com

Received: January 16, 2016; Accepted: February 25, 2016; Published: February 29, 2016

CTTTACACCTTAGCGTATTAACAAGGCAA-CAAGATATCTGTCGTGTTCTCGTCTTGGGCA A T G C C C A A A T C G A C T G C A C C G A C C G A A A C G -G C G A C A C T C C T C T T C A T A T T G C A T G C A G A C T G A G A G A T G A G G G C T G T A T C A G A G C T C T -GACTGAAGGAATATCTCCACTCGAGCGTAAGA GAGGGATGGTTCCACAGAATAGAGCAAGTGGGG-T A C A A C A G C T T C C A C A G A A T C T T G A A C T C A G A A A C T T T G A A G G C T A C A C A T G C A T C -C A T A T T G C A G G A T T C G C T T G T A G C G T C G A T C A G T T G G A G T A C C T T G T G C A G C T A G G C G G C -G A C A T A A A T G C C C C G G A T G G A A A G A G C G G A A G G A C C A T T C T C C A C T A C G C T G T A G A G G C -**G G G T G A C T T T T C T C T T T G T C A G T A C C T C A T T G** CGAACTTGGGTGCCAATGTTAATGCGTTGACCTTTGAC-CAGTGCACACCC3'

C7 was not found.

Discussion and conclusion

The sea star *A. rubens*, although considered to be more primitive than lower vertebrates (as trout) seems to have evolved much more sophisticated immune defense mechanisms.

We find much more complement components in the sea star than in trout: 8 out of 9, when compared to mouse. How do we explain these differences between trout and *A. rubens*?

Phylogenetically speaking the sea star could be situated in "an evolutive cul de sac" and might evolved more quickly than rainbow trout, in term of innate immunity.

As for adaptative immunity, rainbow trout is more evolved [5] than

Asterias rubens which presents an "invertebrate primitive antibody" in response to antigenic injury [6].

This review has described a rather rich catalogue of immune factors in sea star and trout that serve as potent molecules in the defense of these animals against environmental threats.

Taken together one cannot come away with any conclusion other than sea stars have developed a very impressive set of mechanisms to deal with environmental threats.

The same logic would apply to an explanation of why the sea star *A.rubens* has evolved the ability to develop innate and adaptative immunity.

Further studies are bound to unravel the mystery and add to the above information to give a clearly picture of the sequence of events.

References

- Leclerc M, Kresdorn N, Rotter B (2013) Evidence of complement genes in the sea-star Asterias rubens. Comparisons with the sea urchin. Immunol Lett 151: 68-70. [Crossref]
- Zerbino DR, Birney E (2008) Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18: 821-829. [Crossref]
- Wang T, Secombes CJ (2003) Complete sequencing and expression of three complement components, C1r, C4 and C1 inhibitor, of the classical activation pathway of the complement system in rainbow trout Oncorhynchus mykiss. *Immunogenetics* 55: 615-628. [Crossref]
- Chondrou MP, Mastellos D, Zarkadis IK (2006) cDNA cloning and phylogenetic analysis of the sixth complement component in rainbow trout. *Mol Immunol* 43: 1080-7. [Crossref]
- Partula S, Schwager J, Timmusk S, Pilström L, Charlemagne J (1996) A second immunoglobulin light chain isotype in the rainbow trout. *Immunogenetics* 45: 44-51. [Crossref]
- Vincent N, Osteras M, Otten P, Leclerc M (2014) A new gene in A. rubens: A sea star Ig kappa gene. Meta Gene 2: 320-322. [Crossref]

Copyright: ©2016 Leclerc M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.