

Hemorrhage under antivitamins K: a location like no other

Amine Salmi*

Department of Medical and Surgical Emergencies, University Hospital Center Mustapha, Algiers, Algeria

Abstract

Accidents with oral anticoagulants have been widely described, but epidural localization remains exceptional, leading to a misleading clinical picture, the main consequence of which is delayed treatment. The objective of this presentation is to bring out the diagnostic difficulties that arise, to draw the clinician's attention to this location based on a questioning quality. The notion of taking anticoagulants in a patient with neck pain unusual with neurological signs (sensory or motor) is very suggestive, then justify a rapid radiological exploration of the spine.

We report the case of a 34-year-old woman with a mechanical prosthesis in the mitral position (St. Jude), on oral anticoagulant acenocoumarol (Sintrom®) who consults for cervicalgia, overdose with antivitamin K. A cervical scan showed limited epidural hematoma between C3-C6.

Introduction

Oral anticoagulants are usually prescribed for various indications. Their wider use has been noted in recent years, probably exposing more frequent hemorrhagic complications. If, typically, the bleeding is digestive, genitourinary or intracranial, the epidural hematoma, exceptional or even underestimated localization, is rarely reported in the literature. It is characterized by its diagnostic problems on the one hand and by the possible evolutionary complications, in case of late management. Spinal imaging, CT scanning or magnetic resonance imaging (MRI) are very interesting diagnostic tools.

Case presentation

A young 34 years old patient, treated chronically with acenocoumarol (Sintrom®) at a dose of 2 mg per day following a mitral valvuloplasty rather 3 months (St. Jude type), admitted for persistent neck pain, continuous, sudden onset, disobedient to banal analgesic. Physical examination revealed a conscious patient, febrile, without deficit signs or sensory disorders, who complained of severe neck pain, and any attempt to mobilize the cervical spine was impossible. Laboratory tests were not anomalies, outside an INR 7. The brain scan was normal. A cervical spinal cord injury was suspected and CT of the cervical spine then concluded, cervical epidural hematoma at extended from C3-C6 (Figure 1).

All data, clinical and radiological, and absence of any notion of trauma, were in favor of a spontaneous cervical extramedullary hematoma complicating anticoagulation. Anticoagulant therapy was stopped, and the therapeutic management was symptomatic corrected INR (INR <1.5), development of a cervical neck brace for analgesic and clinical monitoring, biological and radiological. The clinical outcome was favorable with disappearance of neck pain and cervical spinal MRI performed a month later, revealed a total resorption of the hematoma (Figure 2).

Discussion

The epidural hematomas (EH) pose a diagnostic problem, given the rarity of this complication, and to our knowledge, we report in this observation the third case of spontaneous EH to overdose on vitamin K, after the two cases cited by Hentschel [1].

The headquarters of the spinal hematoma perished predominates to cervicothoraciques regions, highly mobile regions including the clinical signs depends on two essential elements are: the location and the size of the hematoma [2,3].

On the etiological, our patient had no obvious cause for the fall of the TP, and outside any context of trauma, are the causes of EH [4]:

- The coagulopathy and anticoagulant therapy (30%),
- the arterio-venous malformations (4%),
- tumor cases (4%),
- 30-64% of cases, no cause is found

The risk factors are

- Ankylosing spondylitis,
- bone and ligament aberrations

Thus, 7 cases in EH in the child with hemophilia [5] have been identified, although hit is a rare complication (2.2 to 7.8%) it is necessary to think about the pain of acute spinal installation and this even before the onset of neurological deficit, and treatment should be started at the end of the extension limit. Thrombolytic [6] may be responsible for this complication and 12 cases were reported, with an installation period, neurological disorders, averaged 12 hours.

Treatment [7] of EH can be surgical or conservative, depending on the clinical examination, and are considered good prognostic factors, the following:

Correspondence to: Amine Salmi, Department of Medical and Surgical Emergencies, University Hospital Center Mustapha, Algiers, Algeria, Tel: 0663503399; E-mail: salmi86amine@gmail.com

Key words: epidural hematoma, vitamin K antagonists, misleading location, diagnostic and therapeutic emergency

Received: December 10, 2017; **Accepted:** December 23, 2017; **Published:** December 26, 2017




Figure 1. Cervical epidural hematoma at extended from C3-C6




Figure 2. Total resorption of the hematoma

- a short time between the installation unrest and surgical time (less than 8 hours),
- low compression force,
- a partial deficiency before surgery

Conservative treatment [8] has been tried, and 7 cases of epidural hematoma, which resolved were identified. In our case, the treatment

was based on the rapid antagonism of vitamin K antagonists to obtain an INR (International Normalized Ratio) less than 1.5 associated with analgesic treatment. The patient was referred to her treating physician for the reintroduction of her anticoagulant treatment. In these situations, it is often possible to restart the antivitamin K 2 weeks after the haemorrhagic event.

The prognosis of EH depends on four essential elements [9]:

- the presence of neurological disorders,
- the severity of neurological disorders,
- rapid diagnosis,
- and speed of the treatment.

Evolutionarily, healing is possible:

In total, the deficit is incomplete in 83% of cases where as it is complete in 25% of cases. [10].

Conclusion

The perimedullary hematoma, despite its rarity as a current practice, needs special attention, since the prognosis is directly related to the therapeutic precocity. Careful interrogation and imaging are the diagnostic pillars.

References

1. Hentschel SJ, Woolfenden AR, Fairholm DJ (2001) Resolution of spontaneous spinal epidural hematoma without surgery: report of two cases. *Spine (Phila Pa 1976)* 26: E525–E527. [\[Crossref\]](#)
2. Beatty RM, Winston KR (1984) Spontaneous cervical epidural hematoma: a consideration of etiology. *J Neurosurg* 61: 143–8. [\[Crossref\]](#)
3. Lin IY, Chong C (2004) Non traumatic spinal epidural hematoma mimicking a brainstem stroke. *Ann Emerg Med* 44: 183–184. [\[Crossref\]](#)
4. Domenicucci M, Ramieri A, Ciappetta P, Delfini R (1999) Nontraumatic acute spinal subdural hematoma: report of five cases and review of the literature. *J Neurosurg* 91: 65–73. [\[Crossref\]](#)
5. Chrétiennot-Bara C, Guét A, Balzamo E, Noseda G, Torchet MF, et al. (2001) [Epidural hematoma in a child with hemophilia: diagnostic difficulties]. *Arch Pediatr* 8: 828–833. [\[Crossref\]](#)
6. Cohen J, Ginsberg H, Emery D, Schwartz M (1998) Fatal spontaneous spinal epidural hematoma following thrombolysis for myocardial infarction. *Surg Neurol* 49: 520–523. [\[Crossref\]](#)
7. Groen RJ, van Alphen HA (1996) Operative treatment of spontaneous spinal epidural hematomas: A study of the factors determining postoperative outcome. *Neurosurgery* 39: 494–508. [\[Crossref\]](#)
8. Hentschel SJ, Woolfenden AR, Fairholm DJ (2001) Resolution of spontaneous spinal epidural hematoma without surgery: report of two cases. *Spine (Phila Pa 1976)* 26: E525–527. [\[Crossref\]](#)
9. Otsu I, Merrill D (2003) Epidural hematoma after epidural anesthesia: a case report of non-surgical management. *Acute Pain* 4: 117–120.
10. Cullen D, Bogdanov E, Hüt N (2004) Spinal epidural hematoma occurrence in the absence of known risk factors: a case series. *J Clin Anesth* 16: 376–381. [\[Crossref\]](#)

Copyright: ©2017 Salmi A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.