

SURVEY (Supplemental Information)

Supplemental Information will be excluded from this publication and accessible through a hyperlink

Table 2. Frequencies for Significant Survey Questions (Supplemental Information).

PPGP Risk factors	No. of Pelvic Health PT (%)	No. of Orthopaedic PT (%)	Total (%)
Pain aggravated with prolonged sitting/ standing	30 (97%)	13 (100%)	43 (98%)
Patient points out exact location of pain on body	25 (81%)	13 (100%)	38 (86%)
Patient indicates painful area on diagram	6 (19%)	5 (38%)	11 (25%)
Patient has a history of low back pain	28 (90%)	11 (85%)	39 (89%)
Patient's age	10 (32%)	9 (69%)	19 (62%)
Patient's previous use of contraceptive pills	5 (16%)	2 (15%)	7 (16%)
PPGP Management	No. of Pelvic Health PT (%)	No. of Orthopaedic PT (%)	Total (%)
Stabilization exercise	28 (90%)	13 (100%)	41 (93%)
Motor control exercise	27 (61%)	13 (100%)	40 (90%)
Massage	18 (58%)	7 (54%)	25 (57%)
Acupuncture	4 (13%)	1 (8%)	5 (11%)
Manipulation or joint mobilization	21 (68%)	6 (46%)	27 (61%)
Knowledge of PPGP CPGs	No. of Pelvic Health PT (%)	No. of Orthopaedic PT (%)	Total (%)
Yes	23 (74%)	5 (38.5%)	28 (64%)
No	8 (26%)	8 (61.5%)	16 (36%)
Nonuse of stability/instability terms	No. of Pelvic Health PT (%)	No. of Orthopaedic PT (%)	Total (%)
Agree	25 (81%)	10 (77%)	35 (80%)
Disagree	6 (18%)	3 (23%)	9 (20%)

Copyright: ©2018 Dufour S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.